

Brief Note

Addendum to: Qualitative aspects of solutions in resonators

R. QUINTANILLA¹⁾, R. RACKE²⁾

¹⁾*Department of Applied Mathematics II
UPC Terrassa, Colom 11, 08222 Terrassa, Spain
e-mail: ramon.quintanilla@upc.edu*

²⁾*Department of Mathematics and Statistics
University of Konstanz
78457 Konstanz, Germany
e-mail: reinhard.racke@uni-konstanz.de*

IN CORRECTING A SMALL MISTAKE IN [10], we can prove a new result on non-exponential stability for a coupled system arising in resonators. This also gives another (surprising and simple) example for a thermoelastic system changing from exponential stability to non-exponential stability, when changing from Fourier's law to Cattaneo's law in modeling of the heat conduction.

Key words: hyperbolic model in thermoelasticity, well-posedness, exponential stability, Fourier law, Cattaneo law.

Copyright © 2011 by IPPT PAN

1. Introduction

IN OUR PAPER [10], IN PARTICULAR, the thermoelastic system

$$(1.1) \quad a\Delta^2 u + \Delta\theta + \ddot{u} = 0,$$

$$(1.2) \quad \Delta\theta - m\theta + d\Delta\dot{u} = c\dot{\theta},$$

where

$$(1.3) \quad \hat{f} = f + \tau\dot{f}, \quad \left(\dot{f} = f_t = \frac{\partial f}{\partial t} \right),$$

for the functions $(u, \theta) = (u, \theta)(\mathbf{x}, t)$ with $\mathbf{x} \in B \subset \mathbb{R}^n$, $t \geq 0$, was studied, where B is (smoothly) bounded, $n \geq 2$, and a, m, d, c are positive constants.

¹⁾Research of the first author supported by the project "Partial Differential in Thermomechanics. Theory and Applications (MTM2009-08150).

²⁾Research of the second author supported by the DFG-project "Hyperbolic Thermoelasticity" (RA 504/3-3).

Additionally, one has the boundary conditions

$$(1.4) \quad u(\mathbf{x}, t) = \Delta u(\mathbf{x}, t) = \theta(\mathbf{x}, t) = 0, \quad (\mathbf{x}, t) \in \partial B \times [0, \infty),$$

and initial conditions for $(u, u_t, \theta, \theta_t)$ in $t = 0$. For the background of the interpretation in micro-beam resonators, see [10].

The system is reformulated as a first-order system for

$$\mathbf{V} := (\hat{u}, \hat{u}_t, \theta, \theta_t)',$$

and we obtain

$$(1.5) \quad \mathbf{V}_t = A\mathbf{V}, \quad V(0) = V^0,$$

with the (yet formal) differential operator A given by the symbol:

$$A_f := \begin{pmatrix} 0 & 1 & 0 & 0 \\ -a\Delta^2 & 0 & -\Delta & -\tau\Delta \\ 0 & 0 & 0 & 1 \\ 0 & \frac{d}{c\tau}\Delta & \frac{1}{c\tau}(\Delta - m) & -\frac{1}{\tau} \end{pmatrix}$$

and the initial value

$$\mathbf{V}^0(\mathbf{x}) := (\hat{u}, \hat{u}_t, \theta, \theta_t)'(\mathbf{x}, 0),$$

with its components being given in terms of the originally prescribed initial data by using the differential equations. As an underlying Hilbert space, we have

$$\mathcal{H} := (H^2(B) \cap H_0^1(B)) \times (L^2(B))^n \times H_0^1(B) \times L^2(B)$$

with inner product

$$\begin{aligned} \langle V, W \rangle_{\mathcal{H}} &:= d\langle V^2, W^2 \rangle + ad\langle \Delta V^1, \Delta W^1 \rangle \\ &\quad + \tau\langle \nabla V^3, \nabla W^3 \rangle + \tau m\langle V^3, W^3 \rangle + c\langle V^3 + \tau V^4, W^3 + \tau W^4 \rangle, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the usual $L^2(B)$ -inner product. The operator A is now given as

$$A : D(A) \subset \mathcal{H} \mapsto \mathcal{H}, \quad AV := A_f V,$$

with

$$D(A) := \{V \in \mathcal{H} \mid V^2 \in H^2(B) \cap H_0^1(B), V^4 \in H_0^1(B), A_f V \in \mathcal{H}\}.$$

A generates a contraction semigroup (cf. [10] for the boundary condition $u \in H_0^2(B)$), and A^{-1} is compact, hence the spectrum $\sigma(A)$, of A equals the point spectrum $\sigma_p(A)$.

Due to the boundary conditions, the following Ansatz is possible (cf. [10]):

$$V(t, x) = \sum_{j=1}^{\infty} (\alpha_j(t), \gamma_j(t), \delta_j(t), \varepsilon_j(t))^T w_j(x),$$

for $V = (V^1, V^2, V^3, V^4)^T$, where $(w_j)_j$ denote the eigenfunctions of the Laplace operator under Dirichlet boundary conditions corresponding to the eigenvalue λ_j ,

$$(1.6) \quad -\Delta w_j = \lambda_j w_j, \quad w_j = 0 \quad \text{on} \quad \partial B,$$

with

$$0 < \lambda_1 \leq \dots \leq \lambda_j \rightarrow \infty \quad (\text{as } j \rightarrow \infty).$$

Then the coefficients $\alpha_j, \gamma_j, \delta_j, \varepsilon_j$ satisfy the same differential equation:

$$(1.7) \quad c\tau z'''' + cz''' + (\lambda_j + m + ac\tau\lambda_j^2 + d\tau\lambda_j^2)z'' + (ac\lambda_j^2 + d\lambda_j^2)z' + a\lambda_j^2(\lambda_j + m)z = 0.$$

The corresponding characteristic polynomial P_j is given by

$$(1.8) \quad P_j(\beta) = \beta^4 + \frac{1}{\tau}\beta^3 + \frac{1}{c\tau}(\lambda_j + m + \tau(ac + d)\lambda_j^2)\beta^2 + \frac{1}{c\tau}(ac + d)\lambda_j^2\beta + \frac{a}{c\tau}(\lambda_j^3 + m\lambda_j^2).$$

The zeros of P_j are denoted by $\beta_1(j), \dots, \beta_4(j)$. Let \mathcal{S} denote the spectral set of all zeros,

$$\mathcal{S} := \{\beta_k(j) \mid j = 1, 2, 3, \dots; k = 1, 2, 3, 4\}.$$

Then it is easy to see ([10]) that

$$(1.9) \quad \sigma_p(A) \subset \mathcal{S}.$$

In [10, Theorem 4] it was claimed that

$$(1.10) \quad \exists \omega > 0 : \quad \sup \{\operatorname{Re} \beta \mid \beta \in \mathcal{S}\} \leq -\omega.$$

The final arguments in the proof of (1.10) are based on [10, (5.17)], saying

$$(1.11) \quad \operatorname{Re} \beta_2(j) = \operatorname{Re} \beta_1(j) \longrightarrow -\frac{1}{2\tau},$$

as $j \rightarrow \infty$, which is claimed to follow from [10, (5.8)], saying that

$$(1.12) \quad \operatorname{Re} \beta_1(j) + \operatorname{Re} \beta_3(j) = -\frac{1}{2\tau},$$

and [10, (5.17)], saying that

$$(1.13) \quad \operatorname{Re} \beta_4(j) = \operatorname{Re} \beta_3(j) \longrightarrow -\frac{1}{2\tau}.$$

But (1.12) and (1.13) do *not* imply (1.11), but instead

$$(1.14) \quad \operatorname{Re} \beta_2(j) = \operatorname{Re} \beta_1(j) \longrightarrow 0.$$

This way, the assertion (1.10) (expressing [10, Theorem 4]) is seen to be wrong. But now, the correct relation (1.14) allows us first to prove that the semigroup is *not* exponentially stable (Theorem 1 below).

Since the considerations allow us to take the parameter m in the Eq. (1.2) for θ to be zero without changing the conclusions, we can conclude another new result that gives a new, simple, somehow surprising example for a thermoelastic system, the corresponding semigroup of which changes from exponentially decaying type to non-exponential decaying type, if the model for heat conduction is changed from Fourier's law to Cattaneo's law (Theorem 2 below). This kind of behavior was observed for the rather special Timoshenko system with heat conduction in [2]; now we have another, simpler and even more convincing example that the change from Fourier's law to Cattaneo's law – the modification of a partially parabolic system with infinite propagation speed of signals to a hyperbolic model with finite propagation speed – may change basic properties of the system and hence suggests to think about the modeling character of each system.

2. The non-exponential stability of the semigroup

THEOREM 1.

- (i) $\sup \{\operatorname{Re} \beta \mid \beta \in \mathcal{S}\} = 0$.
- (ii) $\mathcal{S} = \sigma(A) = \sigma_p(A)$.
- (iii) $\{e^{tA}\}_{t \geq 0}$ is not exponentially stable.

P r o o f. Since (i) follows from (1.14), and since (iii) is an easy consequence of (i) and (ii), it remains to show, in view of (1.9), that

$$(2.1) \quad \mathcal{S} \subset \sigma_p(A).$$

To prove Eq. (2.1), let $\beta \in \mathcal{S}$, i.e. $\beta = \beta_k(j)$ for some fixed $j \in \mathbb{N}$ and some fixed $k \in \{1, 2, 3, 4\}$, in particular

$$(2.2) \quad P_j(\beta) = 0.$$

Let $-\Delta w_j = \lambda w_j$, as given in Eq. (1.6). Let $V^1 \in H^2(B) \cap H_0^1(B)$ be the solution of

$$(2.3) \quad \left(\frac{d\beta}{c\tau}\Delta\right)V^1 = \left(\beta^2 + \frac{\beta}{\tau}\right)w_j - \frac{1}{c\tau}(\Delta - m)w_j,$$

and let

$$V := (V^1, \beta V^1, w_j, \beta w_j)^T.$$

Then we have $V \in D(A)$ and we claim:

$$(2.4) \quad AV = \beta V,$$

which would finish the proof. Now, the following equivalences hold: in view of the definition of V^1 in (2.3), the claim (2.4) is equivalent to

$$(2.5) \quad -a\Delta^2 V^1 - \Delta w_j - \tau\beta\Delta w_j = \beta^2 V^1.$$

Applying $\left(\frac{d\beta}{c\tau}\Delta\right)$ to both sides of (2.5), this is equivalent to

$$(2.6) \quad -a\Delta^2\left(\frac{d\beta}{c\tau}\Delta\right)V^1 - \left(\frac{d\beta}{c\tau}\Delta\right)\Delta w_j - \tau\beta\left(\frac{d\beta}{c\tau}\Delta\right)\Delta w_j = \beta^2\left(\frac{d\beta}{c\tau}\Delta\right)V^1,$$

which, using Eqs. (2.3) and (1.6), is equivalent to

$$\left(\beta^4 + \frac{1}{\tau}\beta^3 + \frac{1}{c\tau}(\lambda_j + m + \tau(ac+d)\lambda_j^2)\beta^2 + \frac{1}{c\tau}(ac+d)\lambda_j^2\beta + \frac{a}{c\tau}(\lambda_j^3 + m\lambda_j^2)\right)w_j = 0,$$

which is equivalent to $P_j(\beta) = 0$, and hence it finishes the proof. □

3. Fourier versus Cattaneo law – from exponential to non-exponential stability

As a corollary to the previous Section, we are able to present a new example, where the change from the Fourier law to the Cattaneo law in modeling of the heat conduction, changes the system from an exponentially stable system to a non-exponentially stable one.

This effect was known for a special Timoshenko-type system with heat conduction, and presented in [2]. Here, we give a simpler and even more convincing example for this partially surprising effect that asks for a discussion of the models and their range of validity – even for very simple systems.

In the resonator system discussed above, we may assume $m = 0$ and are still led to the same conclusions. The system – for $m = 0$ from now on – can

be rewritten as follows: introducing a heat flux vector q , a solution (u, θ, q) to

$$(3.1) \quad u_{tt} + a\Delta^2 u + \Delta\theta = 0,$$

$$(3.2) \quad c\theta_t + \operatorname{div} q - d\Delta u_t = 0,$$

$$(3.3) \quad \tau q_t + q + \nabla\theta = 0$$

(plus initial and boundary conditions), gives a solution (u, θ) to the resonator Eqs. (1.1), (1.2) by eliminating q again. The Eq. (3.3) represents the Cattaneo law for heat conduction. As we have seen in the previous section, the solutions do not tend to zero exponentially (and uniformly in the data).

On the other hand, if we take $\tau = 0$ in (3.3) – corresponding to Fourier’s law for heat conduction – we obtain, after eliminating the heat flux q , the classical system for the thermoelastic plate equation:

$$(3.4) \quad u_{tt} + a\Delta^2 u + \Delta\theta = 0,$$

$$(3.5) \quad c\theta_t - \Delta\theta - d\Delta u_t = 0,$$

which is known to be represented by an exponentially stable semigroup, see papers [3, 9, 8, 1, 4, 5, 6, 7]. Hence we have the following

THEOREM 2. *The systems (3.1)–(3.3) and (3.4), (3.5), respectively, give an example where the change from Fourier’s law to Cattaneo’s law in modeling of the heat conduction part, changes the system from an exponentially stable one to a non-exponentially stable one.*

The just presented example for the effect described in the last theorem is simpler than the one presented in [2] which relied on special relations of coefficients, the advantages of which are destroyed by the change from Fourier to Cattaneo.

Acknowledgment

We thank Bugra Kabil for pointing out the mistake in [10] that led to the new results presented here.

References

1. G. AVALOS, I. LASIECKA, *Exponential stability of a thermoelastic system without mechanical dissipation*, Rend. Instit. Mat. Univ. Trieste Suppl., **28**, 1–28, 1997.
2. H. FERNÁNDEZ SARE, R. RACKE, *On the stability of damped Timoshenko systems – Cattaneo versus Fourier law*, Arch. Rational Mech. Anal., **194**, 221–251, 2009.

3. J.U. KIM, *On the energy decay of a linear thermoelastic bar and plate*, SIAM J. Math. Anal., **23**, 889–899, 1992.
4. I. LASIECKA, R. TRIGGIANI, *Two direct proofs on the analyticity of the S.C. semigroup arising in abstract thermoelastic equations*, Adv. Differential Equations, **3**, 387–416, 1998.
5. I. LASIECKA, R. TRIGGIANI, *Analyticity, and lack thereof, of thermo-elastic semigroups*, ESAIM, Proc., **4**, 199–222, 1998.
6. I. Lasiecka, R. Triggiani, *Analyticity of thermo-elastic semigroups with coupled hinged/Neumann boundary conditions*, Abstract Appl. Anal., **3**, 153–169, 1998.
7. I. LASIECKA, R. TRIGGIANI, *Analyticity of thermo-elastic semigroups with free boundary conditions*, Annali Scuola Norm. Sup. Pisa, **27**, 457–482, 1998.
8. Z. LIU, S. ZHENG, *Exponential stability of the Kirchhoff plate with thermal or viscoelastic damping*, Quart. Appl. Math., **53**, 551–564, 1997.
9. J.E. MUÑOZ RIVERA, R. RACKE, *Smoothing properties, decay and global existence of solutions to nonlinear coupled systems of thermoelastic type*, SIAM J. Math. Anal., **26**, 1547–1563, 1995.
10. R. QUINTANILLA, R. RACKE, *Qualitative aspects in resonators*, Arch. Mech., **60**, 345–360, 2008.

Received January 25, 2011.
